



## The Relationship Between The Level of Knowledge and Family Support, and Measures to Prevent Pulmonary Tuberculosis Transmission

Tessalonika Marcelin Pangaribuan<sup>1</sup>, Ika Mustafida<sup>1\*</sup>, Enni Juliani<sup>1</sup>

<sup>1</sup> Bachelor of Nursing Study Program, STIKES RS Husada, Central Jakarta, DKI Jakarta, Indonesia

### ARTICLE INFO

**Article Type:**  
Research

**Article History:**  
Received: 03 August 2025  
Accepted: 29 January 2026  
Published: 31 March 2026

**\*Corresponding author**  
Email: [ikamustafida@gmail.com](mailto:ikamustafida@gmail.com)

### ORIGINAL ARTICLE

#### ABSTRACT

Pulmonary tuberculosis remains a global health problem with a significant burden, particularly in densely populated areas such as Jakarta. The occurrence of this disease is influenced not only by medical factors but also by the level of individual knowledge and family support in preventing its transmission. This study aims to analyze the relationship between knowledge level and family support with efforts to prevent the transmission of pulmonary tuberculosis in the working area of the Kemayoran Public Health Center, Central Jakarta. This study employed a quantitative approach with a cross-sectional design involving 75 respondents selected through a survey technique using a structured questionnaire. Data analysis was conducted using univariate analysis to describe variable distributions and bivariate analysis using Spearman's rho correlation test. The results showed a significant relationship between knowledge level and efforts to prevent the transmission of pulmonary tuberculosis, with a strong correlation. In addition, family support also demonstrated a very strong and significant relationship with prevention efforts. The majority of respondents had a moderate level of knowledge, tended to have low family support, and their prevention efforts were generally in the moderate to good category. In conclusion, this study highlights that improving knowledge and strengthening family support play an important role in enhancing preventive behaviors against pulmonary tuberculosis transmission. Therefore, continuous health education interventions and family empowerment are needed to reduce the transmission rate of pulmonary tuberculosis, particularly in urban areas.

**Keywords:** Pulmonary Tuberculosis, Knowledge Level, Family Support, Prevention Efforts.

#### ABSTRAK

Tuberkulosis paru masih menjadi masalah kesehatan global dengan beban yang signifikan, terutama di wilayah padat penduduk seperti Jakarta. Kejadian penyakit ini tidak hanya dipengaruhi oleh faktor medis, tetapi juga berkaitan dengan tingkat pengetahuan individu serta dukungan keluarga dalam upaya pencegahan penularan. Penelitian ini bertujuan untuk menganalisis hubungan antara tingkat pengetahuan dan dukungan keluarga dengan upaya pencegahan penularan tuberkulosis paru di wilayah kerja Puskesmas Kemayoran, Jakarta Pusat. Penelitian ini menggunakan pendekatan kuantitatif dengan desain cross-sectional terhadap 75 responden yang dipilih melalui teknik survei menggunakan kuesioner terstruktur. Analisis data dilakukan secara univariat untuk menggambarkan distribusi variabel dan secara bivariat menggunakan uji korelasi Spearman's rho. Hasil penelitian menunjukkan adanya hubungan yang signifikan antara tingkat pengetahuan dengan upaya pencegahan penularan tuberkulosis paru dengan kekuatan hubungan yang kuat. Selain itu, dukungan keluarga juga memiliki hubungan yang sangat kuat dan signifikan dengan upaya pencegahan penularan tuberkulosis paru. Mayoritas responden memiliki tingkat pengetahuan dalam kategori cukup, dukungan keluarga cenderung rendah, serta upaya pencegahan penularan berada pada kategori cukup hingga baik. Kesimpulan penelitian ini menegaskan bahwa peningkatan pengetahuan dan penguatan dukungan keluarga berperan penting dalam meningkatkan perilaku pencegahan penularan tuberkulosis paru. Oleh karena itu, diperlukan intervensi berupa edukasi kesehatan yang berkelanjutan serta pemberdayaan keluarga guna menekan angka penularan tuberkulosis paru, khususnya di wilayah perkotaan.

**Kata Kunci:** TB Paru, Tingkat Pengetahuan, Dukungan Keluarga, Upaya Pencegahan Penularan.

## **INTRODUCTION**

Pulmonary tuberculosis is an infectious disease caused by *Mycobacterium tuberculosis* and remains a global health problem with a significant burden, particularly in developing countries including Indonesia. The disease can infect one-third of the world's population and is among the top ten leading causes of death globally, with an estimated 10.6 million cases annually (Sibua, 2021; Kementerian Kesehatan Republik Indonesia, 2022). In Indonesia, the prevalence of tuberculosis reaches 9.2%, with a higher proportion in males compared to females, at 57.9% and 42.1%, respectively (Kementerian Kesehatan Republik Indonesia, 2022). The risk factors for tuberculosis are not only related to medical aspects but are also influenced by sociodemographic factors such as age, education, occupation, and income, as well as environmental and behavioral factors including housing conditions, smoking habits, and comorbidities (Pralambang & Setiawan, 2021). Airborne droplet transmission makes this disease highly contagious, particularly in densely populated areas.

Clinically, pulmonary tuberculosis is characterized by symptoms such as persistent productive cough, hemoptysis, chest pain, fever, weight loss, and night sweats (Rokhmah, 2013). If not properly treated, the disease can progress into serious complications, both in early and advanced stages, including pleural effusion, lung parenchymal damage, acute respiratory distress syndrome (ARDS), and death due to massive hemoptysis (Pratiwi, 2020). Beyond its clinical impact, pulmonary tuberculosis also affects patients' quality of life, including physical, psychological, and social aspects (Nainggolan, 2022). Therefore, efforts to prevent transmission are a crucial component in controlling this disease, particularly within families and communities.

Efforts to prevent the transmission of pulmonary tuberculosis are strongly influenced by individual knowledge levels and family support. Adequate knowledge regarding the disease, its transmission, and treatment can enhance awareness and promote preventive behaviors (Lutfian, 2025). Meanwhile, family support, both emotional and instrumental, plays a significant role in improving patient adherence to treatment and healthy lifestyle behaviors. However, previous studies have shown inconsistent findings. Research by Hayati and Sulistiyowati (2024) indicated no significant relationship between knowledge level and preventive behavior, while a significant association was found between family support and tuberculosis prevention behavior ( $p$ -value = 0.009). These differing findings highlight a research gap that requires further investigation.

Epidemiologically, the DKI Jakarta region continues to experience a high burden of tuberculosis cases, with a total of 33,552 cases, of which Central Jakarta accounts for 5,008 cases (Badan Pusat Statistik Provinsi DKI Jakarta, 2021). Data from the Kemayoran Public Health Center show a significant increase in cases from 83 cases in 2023 to 303 cases in 2024. Preliminary study findings indicate low levels of patient knowledge regarding tuberculosis and insufficient family support, influenced by educational and economic factors. Additionally, high population density and unhealthy lifestyle behaviors such as smoking further exacerbate the risk of transmission. These conditions suggest that behavioral and social factors play a critical role in the increasing incidence of pulmonary tuberculosis in this area.

Although numerous studies have examined factors influencing pulmonary tuberculosis, most have focused on treatment adherence and therapeutic outcomes. Studies specifically analyzing the relationship between knowledge level and family support with transmission prevention efforts, particularly in densely populated urban settings such as Central Jakarta, remain limited. Furthermore, inconsistencies in previous findings indicate an empirical gap that needs to be addressed. Therefore, the novelty of this study lies in the simultaneous analysis of knowledge and family support factors in relation to tuberculosis transmission prevention efforts within a high-density urban community context. This study aims to analyze the relationship between knowledge level and family support with efforts to prevent the transmission of pulmonary tuberculosis in the working area of the Kemayoran Public Health Center, Central Jakarta.

## **RESEARCH METHODS**

This study is a quantitative study with a cross-sectional design aimed at analyzing the relationship between knowledge level and family support with efforts to prevent the transmission of pulmonary tuberculosis. The population in this study consisted of all pulmonary tuberculosis patients who were recorded and undergoing treatment in the working area of the Kemayoran

Public Health Center, Central Jakarta, totaling 75 individuals. The sampling technique used was total sampling; therefore, all members of the population who met the inclusion and exclusion criteria were included as research respondents.

The inclusion criteria in this study were pulmonary tuberculosis patients registered at the Kemayoran Public Health Center, aged  $\geq 18$  years, and willing to participate as respondents by providing informed consent. Meanwhile, the exclusion criteria included patients with physical or mental conditions that made them unable to complete the questionnaire independently.

Data were collected using a structured questionnaire that had undergone validity and reliability testing to ensure the appropriateness of the instrument in measuring the research variables. The variables examined included knowledge level, family support, and efforts to prevent the transmission of pulmonary tuberculosis.

Data analysis was conducted using univariate analysis to describe the frequency distribution and characteristics of each research variable. Furthermore, bivariate analysis was performed using Spearman's rho correlation test to identify the relationship between knowledge level and family support with efforts to prevent the transmission of pulmonary tuberculosis. This study obtained ethical approval (ethical clearance) from the Health Research Ethics Committee with approval number No.1122/KEPK-FIK/V/2025 prior to data collection, ensuring that all research procedures complied with applicable ethical research principles.

## RESULTS

**Table 1.** Distribution of Respondent Characteristics Based on Age, Education Level, Occupation, and Gender among Pulmonary Tuberculosis Patients at Kemayoran Public Health Center (n = 75).

Variable	Category	Frequency (n)	Percentage (%)
Age	$\leq 25$ years	42	56.0
	$> 25$ years	33	44.0
	Total	75	100
Education	No formal education	8	10.7
	Primary school	4	5.3
	Junior high school	11	14.7
	Senior high school	29	38.7
	Higher education	23	30.7
	Total	75	100
Occupation	Unemployed	18	24.0
	Government/Army/Police	18	24.0
	Housewife	13	17.3
	Self-employed	15	20.0
	Others	11	14.7
	Total	75	100
Gender	Male	32	42.7
	Female	43	57.3
	Total	75	100

Based on Table 1, the majority of respondents were aged  $\leq 25$  years (56.0%), indicating that most participants were in the productive age group, which is generally associated with higher mobility and increased exposure to infectious diseases such as pulmonary tuberculosis. In terms of education, most respondents had completed senior high school (38.7%), followed by higher education (30.7%), while a smaller proportion had only basic education or no formal education. This variation suggests that educational background may influence health literacy and awareness regarding disease prevention. Regarding occupation, the largest proportions were government employees (24.0%) and unemployed individuals (24.0%), followed by self-employed workers (20.0%), indicating diverse socioeconomic conditions that may affect exposure risk and access to healthcare. In terms of gender, most respondents were female (57.3%), which may

reflect differences in healthcare-seeking behavior or exposure patterns in densely populated environments.

**Table 2.** Distribution of Knowledge Level among Respondents (n = 75).

Knowledge Level	Frequency (n)	Percentage (%)
Poor	24	32.0
Moderate	27	36.0
Good	24	32.0
Total	75	100

Table 2 shows that the majority of respondents had a moderate level of knowledge (36.0%), while equal proportions were observed in the poor and good categories (32.0% each). This indicates that although some respondents have adequate knowledge, a considerable proportion still lacks sufficient understanding of tuberculosis and its prevention.

**Table 3.** Distribution of Family Support among Respondents (n = 75).

Family Support	Frequency (n)	Percentage (%)
Very Low	15	20.0
Low	23	30.7
Moderate	13	17.3
High	12	16.0
Very High	12	16.0
Total	75	100

Table 3 indicates that most respondents reported low family support (30.7%), followed by very low support (20.0%). Only a small proportion experienced high or very high support. This finding suggests that family involvement in patient care and prevention efforts remains suboptimal, which may negatively influence treatment adherence and preventive behaviors.

**Table 4.** Distribution of Tuberculosis Transmission Prevention Efforts (n = 75).

Prevention Efforts	Frequency (n)	Percentage (%)
Poor	24	32.0
Moderate	27	36.0
Good	24	32.0
Total	75	100

Table 4 reveals that the majority of respondents demonstrated moderate prevention efforts (36.0%), while equal proportions were categorized as poor and good (32.0% each). This distribution indicates that preventive practices are not yet optimal and still vary widely among patients.

**Table 5.** Bivariate Analysis of the Relationship between Knowledge Level, Family Support, and Tuberculosis Transmission Prevention Efforts (n = 75)

Variables	Correlation Coefficient (r)	p-value	Interpretation
Knowledge Level – Prevention Efforts	0.682	0.001	Strong correlation
Family Support – Prevention Efforts	0.855	0.001	Very strong correlation

Table 5 shows the results of the bivariate analysis using Spearman's rho correlation test indicate that there is a statistically significant relationship between knowledge level and tuberculosis transmission prevention efforts ( $p = 0.001$ ). The correlation coefficient ( $r = 0.682$ ) shows a strong positive relationship, meaning that higher levels of knowledge are associated with better prevention practices among patients with pulmonary tuberculosis.

Furthermore, family support also demonstrates a statistically significant relationship with prevention efforts ( $p = 0.001$ ). The correlation coefficient ( $r = 0.855$ ) indicates a very strong

positive relationship, suggesting that patients who receive higher levels of family support are more likely to engage in effective tuberculosis transmission prevention behaviors.

## **DISCUSSION**

The characteristics of respondents in this study indicate that the majority were in the productive age group. This condition can be explained by the fact that individuals in this age group tend to have high levels of activity and mobility, thereby increasing their risk of exposure to *Mycobacterium tuberculosis*. In addition, work demands and intensive social activities may reduce immune function due to fatigue and lack of rest. These findings are consistent with previous studies indicating that the productive age group has a higher risk of pulmonary tuberculosis compared to other age groups (Sari et al., 2019; Susanti et al., 2024; Lauchan et al., 2025). Therefore, this age group should be a primary target for tuberculosis prevention programs, particularly through active screening and workplace- and community-based health education.

In terms of education, the majority of respondents had a secondary education level (senior high school). Education plays an important role in shaping an individual's ability to understand health-related information, including tuberculosis transmission and prevention. However, formal education does not always directly translate into preventive behavior if it is not supported by continuous and contextual health education. This finding is consistent with studies by Berhan et al. (2023) and Firmanda et al. (2025), which reported that high school graduates dominate tuberculosis patients but still require strengthened health education. This highlights that improving health literacy depends not only on formal education but also on sustained educational interventions.

Based on occupation, most respondents were categorized as unemployed. This condition is associated with economic limitations that may affect access to health information, healthcare services, and adequate living environments. Unemployed individuals also tend to spend more time in densely populated home environments, thereby increasing the risk of tuberculosis transmission (Andriani, Lestari, & Prasida, 2023). This finding is in line with studies by Hasnita et al. (2025) and Umar (2021), which indicate that unemployment is associated with an increased risk of pulmonary tuberculosis. However, it should be critically noted that employment status does not act independently but interacts with other factors such as education, housing conditions, and access to healthcare services.

In terms of gender, the majority of respondents were female. This reflects the social role of women within the family as primary caregivers and managers of household health, which increases their likelihood of exposure to infectious diseases such as tuberculosis. Additionally, women tend to be more proactive in accessing healthcare services compared to men. These findings are consistent with studies by Wulandari et al. (2020) and Guido et al. (2024). However, the predominance of female respondents may also indicate that men are more likely to delay seeking healthcare, potentially leading to underdiagnosis among male patients. Therefore, a gender-based approach should be considered in tuberculosis control programs.

The level of knowledge among respondents was predominantly in the moderate category, indicating that information about tuberculosis has been received but not fully understood or optimally applied. Incomplete knowledge may limit individuals' ability to identify transmission risks and consistently implement preventive behaviors. This finding aligns with studies by Fikri, Pelawi, & Deniati (2024), which highlight the gap between information exposure and public understanding of tuberculosis. This supports health behavior theory, which suggests that knowledge is a prerequisite but not sufficient to drive behavioral change without the support of other factors.

Family support in this study varied, with a tendency toward low to moderate levels. This indicates that family involvement in supporting tuberculosis patients is not yet optimal. In fact, family support, both emotional and instrumental, plays a crucial role in improving treatment adherence and the implementation of preventive behaviors. These findings are consistent with studies by Fawwaz (2022), which emphasize the importance of family support in tuberculosis control. Theoretically, this finding reinforces the social ecological approach, which positions the family as a key factor in shaping individual health behaviors (Sari, 2023).

The prevention efforts for tuberculosis transmission in this study were mostly categorized as moderate, indicating that preventive practices have been implemented but are not yet optimal. This suggests the presence of barriers to consistently applying preventive behaviors, despite the availability of information. Factors such as stigma, lifestyle habits, and densely populated living conditions are major obstacles to tuberculosis prevention (Darliana, 2011). Therefore, prevention strategies should adopt a comprehensive approach that considers individual, social, and environmental factors.

The results of the bivariate analysis showed a significant relationship between knowledge level and family support with tuberculosis prevention efforts. The correlation between knowledge level and prevention efforts was categorized as strong, while the relationship between family support and prevention efforts was categorized as very strong. These findings indicate that individual and social factors complement each other in shaping preventive behaviors. Notably, family support was found to have a stronger influence than knowledge level in improving prevention efforts. This suggests that interventions focusing solely on increasing knowledge may not be sufficient without involving the immediate social environment, particularly the family.

These findings highlight that both knowledge level and family support play important roles in improving tuberculosis prevention efforts. However, family support appears to have a stronger influence compared to knowledge level. This emphasizes the importance of involving family members in tuberculosis control programs, alongside enhancing patient education to promote optimal preventive behaviors. Therefore, tuberculosis control programs should be designed holistically by integrating health education and family empowerment as a unified intervention strategy.

This study has several limitations that should be considered when interpreting the findings. First, the study was conducted only in the Kemayoran Public Health Center area, limiting the generalizability of the results to other regions with different characteristics. Second, the cross-sectional design does not allow for the establishment of causal relationships. Third, data collection using questionnaires may introduce subjective bias from respondents. Therefore, future studies are recommended to use longitudinal designs and include additional variables such as environmental and cultural factors to provide a more comprehensive analysis.

## **CONCLUSION**

Based on the results of this study, it can be concluded that knowledge level and family support play a significant role in efforts to prevent the transmission of pulmonary tuberculosis in the working area of the Kemayoran Public Health Center, Central Jakarta. The majority of respondents had a moderate level of knowledge; however, this understanding has not been consistently implemented in daily behavior. On the other hand, family support tended to be low, indicating that family involvement in supporting treatment and disease prevention remains suboptimal, influenced by limited knowledge, economic conditions, and persistent social stigma within the community.

Nevertheless, respondents demonstrated prevention efforts at a moderate to good level, such as maintaining proper home ventilation, practicing cough etiquette, and seeking health information. The results of the bivariate analysis showed a significant relationship between knowledge level and family support with prevention efforts, with family support exhibiting a stronger correlation compared to knowledge level.

Therefore, comprehensive interventions are needed, not only focusing on improving individual knowledge through continuous health education but also on strengthening the role of the family as the primary support system for patients. Family empowerment is expected to enhance the consistency of preventive behaviors, improve treatment adherence, and reduce the transmission rate of pulmonary tuberculosis, particularly in densely populated areas such as Kemayoran.

## **REFERENCES**

Andriani, L., Lestari, R. M., & Prasida, D. W. (2023). Hubungan tingkat pengetahuan serta dukungan keluarga dengan kepatuhan minum obat pada penderita tuberculosis di wilayah

- kerja Puskesmas Pahandut. *Jurnal Surya Medika*, 9(1), 96–103. <https://doi.org/10.33084/jsm.v9i1.5155>
- Badan Pusat Statistik Provinsi DKI Jakarta. (2021). *Provinsi DKI Jakarta Dalam Angka 2021*. Jakarta: Badan Pusat Statistik Provinsi DKI Jakarta.
- Berhan, A., Almagaw, A., Solomon, Y., Legese, B., Damtie, S., Erkihun, M., Alebachew, Z., Eyayu, T., & Getie, G. A. B. (2023). Tuberculosis treatment outcome and associated factors among tuberculosis patients linked to tuberculosis treatment clinics in Ethiopia, 2023: A multi-center retrospective study. *Infection and Drug Resistance*, 16, 3367–3378. <https://doi.org/10.2147/IDR.S413272>
- Darlina, D. (2011). Manajemen pasien tuberculosis paru. *Idea Nursing Journal*, 2(1), 27–31.
- Fawwaz, F., Susanto, A., & Sukmaningtyas, W. (2022). Hubungan tingkat pengetahuan dengan pencegahan penularan tuberculosis paru di Balai Kesehatan Paru Masyarakat (BKPM) Purwokerto. *Jurnal Sehat Mandiri*, 17(2), 69–77. <https://doi.org/10.33761/jsm.v17i2.673>
- Fikri, M., Pelawi, A., & Deniati, K. (2024). Hubungan Tingkat Pengetahuan Pasien TB Paru dengan Upaya Pencegahan Penularan TB Paru. *Jurnal Penelitian Perawat Profesional*, 6(4), 1565-1574. Retrieved from <https://jurnal.globalhealthsciencegroup.com/index.php/JPPP/article/view/2805>
- Firmanda, G. I., Pratiwi, W. N., Sunarno, R. D., & Wahyuningsih, A. (2025). Pengaruh edukasi terhadap pengetahuan dan kepatuhan obat pada penderita TB di Karanganyar. *Jurnal Keperawatan Klinis dan Komunitas (Clinical and Community Nursing Journal)*, 9(1), 28–36. <https://doi.org/10.22146/jkck.104297>
- Guido, R. M., Juniartati, E., Putro, W., & Fittarsih, N. (2024). Hubungan pengetahuan, sikap, dukungan keluarga terhadap perilaku pencegahan penularan tuberculosis di wilayah kerja Puskesmas Ngabang Kabupaten Landak Tahun 2024. *Scientific Journal of Nursing Research*, 6(1), 11-19. Retrieved from <https://ejournal.poltekkes-pontianak.ac.id/index.php/SJNR/article/view/1561>
- Hasnita, N., Ichwansyah, F., Abdurrahman, F. bin, Usman, S., & Zahara, M. (2025). Analysis of Risk Factors for Pulmonary Tuberculosis Incidence. *Indonesian Journal of Global Health Research*, 7(2), 219-234. <https://doi.org/10.37287/ijghr.v7i2.5553>
- Hayati, U., & Sulistiyowati, A. N. (2024). The Influence of Housing Density, History of Contact with Positive TB Cases, And BCG Immunization History on The Occurrence of Pulmonary Tuberculosis In Productive Age Individuals. *Jurnal EduHealth*, 15(04), 1033-1044. Retrieved from <https://ejournal.seaninstitute.or.id/index.php/health/article/view/6034>
- Kementerian Kesehatan Republik Indonesia. (2022). *Profil kesehatan Indonesia 2021*. Kementerian Kesehatan Republik Indonesia.
- Lauchan, A. M., Ashar, Y. K., Siregar, P. A., & Samosir, H. E. (2025). Determinant factors of lung tuberculosis in North Sumatera province: Analysis of Indonesia health survey data 2023. *Gema Lingkungan Kesehatan*, 23(2), 211–220. <https://doi.org/10.36568/gelinkes.v23i2.283>
- Lutfian, L., Azizah, A., Wardika, I. J., Wildana, F., Maulana, S., & Wartakusumah, R. (2025). The role of family support in medication adherence and quality of life among tuberculosis patients: A scoping review. *Japan Journal of Nursing Science*, 22(1), e12629. <https://doi.org/10.1111/jjns.12629>
- Nainggolan, M. (2022). Hubungan Pengetahuan, Sikap dan Dukungan Keluarga terhadap Perilaku Pencegahan Penularan Pada Pasien TBC: Relationship of Knowledge, Attitude, and Family Support to Prevention of Transmission of Behavior in TB Patients. *Indonesian Scholar Journal of Medical and Health Science*, 1(08), 300-305. Retrieved from <https://dohara.or.id/index.php/hsk/article/view/179>
- Pralambang, S. D., & Setiawan, S. (2021). Faktor risiko kejadian tuberculosis di Indonesia. *Jurnal Biostatistik, Kependudukan, serta Informatika Kesehatan*, 2(1), 60-71. <https://doi.org/10.51181/bikfokes.v2i1.4660>
- Pratiwi, R. D. (2020). Gambaran komplikasi penyakit Tuberculosis berdasarkan kode international classification of disease 10. *Jurnal Kesehatan Al-Irsyad*, 13(2), 93-101.

- Rokhmah, D. (2013). Gender dan Penyakit Tuberkulosis: Implikasinya Terhadap Akses Layanan Kesehatan Masyarakat Miskin yang Rendah. *Kesmas*, 7(10), 447-452. <https://doi.org/10.21109/kesmas.v7i10.3>
- Sari, D. K., Mega, J. Y., & Harahap, J. (2019). Nutrition status related to clinical improvement in AFB-positive pulmonary tuberculosis patients in primary health centres in Medan, Indonesia. *Open Access Macedonian Journal of Medical Sciences*, 7(10), 1621–1627. <https://doi.org/10.3889/oamjms.2019.338>
- Sari, M., Efriyandi, E., Tamsil, T. A., & Lestrari, I. C. (2023). Hubungan tingkat pengetahuan penyakit TB dengan pencegahan penularan TB pada masyarakat Desa Meranti. *Ibnu Sina: Jurnal Kedokteran Dan Kesehatan - Fakultas Kedokteran Universitas Islam Sumatera Utara*, 22(1), 6-10. <https://doi.org/10.30743/ibnusina.v22i1.326>
- Sibua, S. (2021). Hubungan dukungan keluarga dengan kepatuhan berobat penderita tuberkulosis di Kabupaten Bolaang Mongondow Timur. *Aksara: Jurnal Ilmu Pendidikan Nonformal*, 7(3), 1443–1450. <https://doi.org/10.37905/aksara.7.3.1443-1450.2021>
- Susanti, E. L., Senen, P., Hasnaini, A., & Senen, P. (2024). Karakteristik dan analisis risiko kasus tuberkulosis di Puskesmas Senen, Jakarta Pusat tahun 2023. *Epidkes*, 8(2). <https://doi.org/10.7454/epidkes.v8i2.1130>
- Umar, E. (2021). Faktor-faktor yang berhubungan kejadian TBC dengan pengetahuan, perilaku, dan lingkungan sosial penderita TBC di RW I Kelurahan Terondol Kota Serang. *Jurnal Ilmiah Keperawatan*, 2(2), 25–32.
- Wulandari, I. S. M., Rantung, J., & Malinti, E. (2020). Faktor yang berhubungan dengan kepatuhan minum obat pasien TB di wilayah kerja Puskesmas Parongpong. *Jurnal Keperawatan Muhammadiyah*, 5(1), 128-134. <https://doi.org/10.30651/jkm.v5i1.4536>